

## **LCLU WP Summer internships Projects**

Please email [admin@lclu.cam.ac.uk](mailto:admin@lclu.cam.ac.uk) with any queries, or contact the supervisors directly about the projects.

### **1. Did magnetite cause biological molecules to become “one-handed”?**

*Supervisors: Prof Nicholas Tosca and Prof Richard Harrison, Department of Earth Sciences*

*Co-applicants: Clancy Jiang, Dept of Earth Sciences; S. Furkan Ozturk and Dimitar Sassellov, Harvard University*

Homochirality, or the exclusive selection of one of two possible mirror-image forms of sugars, amino acids, nucleotides and their respective polymers, is a fundamental signature of all life on Earth. However, the origins of homochirality have persisted as a central problem in research related to life's origins. A recent hypothesis has suggested that the mineral magnetite could have provided a source of spin-polarized electrons which may have led to the enrichment of one chiral form of bimolecular building blocks at a critical point in the origins of life. However, the efficiency of this process depends on the physical properties (size, shape, magnetic structure) of magnetite. What environments on the early Earth could have generated magnetite with suitable properties to induce homochirality?

To test the hypothesis that magnetite may have played a key role in the origin of homochirality, the aim of this project will be to investigate prebiotically-relevant synthesis routes for magnetite on the early Earth, and how its physical and magnetic properties vary under different conditions. The methodology will involve magnetite synthesis from oxygen-free water under a range of conditions relevant to prebiotic environments using experimental facilities in the Aqueous Geochemistry Laboratory, and a range of materials characterization facilities in the Department of Earth Sciences. The magnetic properties of the synthetic magnetite will be characterized in the NanoPalaeomagnetism Laboratory, and the results will be synthesized in the context of current constraints on the efficiency of inducing enantiomeric excess with spin-polarized electrons. We anticipate the results will carry important implications for the natural environments and conditions under which magnetite may have facilitated enantiomeric excess on the early Earth, providing new perspective on the origins of life.

### **2. Can you detect an Exoplanet for <£100?**

*Clark Baker, Matthew Hooton, Cavendish Astrophysics*

In this project, the student will investigate whether you can detect an exoplanet transit from Cambridge using equipment that costs £100 or less.

Initially, the student will be given a crash course in observational astronomy. Using this knowledge, they will build an exposure time calculator (ETC)/ observation

simulator for the photometric observation of an exoplanet. This will give the intern a good grounding and understanding of observational exoplanet science.

Applying the tool that the student has built, they will investigate the number of exoplanet transits from the exoplanet archive that they could re-observe with a small (0.1-0.3m) aperture telescope.

Time permitting, the student will look to procure the best instrument they can (pre-owned telescope and detector) with their modest budget of ~£100 and attempt to make one of the observations that they have simulated.

The lead applicant specialises in instrumentation for exoplanet science, is a member of the HARPS3 team and has previously developed a spectroscopic ETC and extremely low-cost radial velocity spectrograph.

### **3. How to make a fossil – and interpret the results**

*Supervisors: Prof Nick Butterfield, Department of Earth Sciences*

Fossils provide the only direct record of the deep-time history of life on Earth, but the pathways to producing them are complex and rarely successful. At the same time, there is any number of abiotic processes that can generate biology-mimicking form – pseudofossils. This project will examine a remarkable assemblage of fossils and pseudofossils from the ~100 million-year-old (mid-Cretaceous) Cambridge Greensand, with an eye to understanding the circumstances responsible for their preservation/formation, and what that might tell us about the biological, environmental and geochemical nature of the Greensand seas.

In addition to being green and sandy, the Cambridge Greensand is famous for its abundance of phosphatic ‘coprolites’ and the basis of the world’s first artificial fertilizer. Although not actually fossil faeces, the industrialized excavation of phosphate-rich horizons in the late 19th century yielded an enormous diversity of bona fide fossils – from single celled phytoplankton to sponges, molluscs, echinoderms, aquatic reptiles and dinosaurs – most of which were similarly preserved via a process of phosphate biomineralization. This project will zero in on a number of these fossils and fossil-like structures in order to understand the sequence of events leading to their formation, not least the interplay between the original biominerals (calcite, aragonite, silica, phosphate), their selective preservation/dissolution/replacement, and the background effects of contemporaneous biological and sedimentological reworking.

The day to day work will involve the production and analysis of multiple petrographic thin-sections of Cambridge Greensand body fossils, trace fossils (burrows and borings) and ‘coprolites.’ Most of the analytical work will be conducted with a petrographic microscope, though there may also be an opportunity for some electron microscopy. Serial sectioning and virtual reassembly of key specimens would also be a possibility. On a more creative level, the student would be encouraged to think

about how the preserved constituents relate back to the once-living organisms and their post-mortem biogeochemical processing.

#### **4: Impact of the first arthropods on the biosphere**

*Supervisors: Dr Stephen Pates and Dr Emily Mitchell, Department of Zoology*

Arthropods have dominated the animal world since they first appeared during the Cambrian Explosion half a billion years ago. They were critical for construction and maintenance of the animal dominated biological pump throughout this time through mixing of the water column, playing crucial roles in the early Palaeozoic carbon cycle just as they do in the modern. However, links between their wide array of morphologies and their hydrodynamic impact have not been quantified.

Computational fluid dynamics approaches to determine the hydrodynamic performance of an array of arthropod morphologies provide an excellent opportunity to link form and function in arthropods, while morphometric and geometric morphometric approaches facilitate comparisons between extant and extinct forms. For this research project, specimens of modern crustaceans in the Cambridge Museum of Zoology will be utilised to quantify links between form and function in arthropods, providing valuable information that will provide insights into how the early diversification of form in Cambrian arthropods facilitated different life modes. Three-dimensional models of specimens of modern crustaceans in the Cambridge Museum of Zoology collections will be constructed. Specimens will be chosen that vary in carapace shape, abdomen length, and other morphological features seen to greatly vary in Cambrian arthropods. The hydrodynamic performance of these models will be quantified using computational fluid dynamics analyses. A range of conditions will be considered, including varying flow speeds and distances from the seafloor, to understand how morphology links to hydrodynamics in different environments and for different life modes. The morphology of extant specimens will then be quantitatively compared to Cambrian arthropods (data gathered from the literature). This will allow inference of how the morphological variation in these Cambrian arthropods impacted their hydrodynamic performance. Thus, the suitability of different Cambrian arthropod morphologies for different modes of life in these early animal ecosystems will be revealed.